



Misoprostol Titration Protocol

Southern Health-Santé Sud Oral Misoprostol Titration Protocol "A" (200 mcg tablet)

Time/Progress Marker	Dose	Route
<i>Dissolve 200 mcg Misoprostol tablet in 10 mL of water = final concentration 20mcg/mL Dilute further in 30 mL of water for patient to ingest full dose</i>		
First dose after admission	20 mcg (1 mL)	Oral
After 2 hours	20 mcg (1 mL)	Oral
After 4 hours	20 mcg (1 mL)	Oral
After 6 hours	30 mcg (1.5 mL)	Oral
After 8 hours	30 mcg (1.5 mL)	Oral
After 10 hours	30 mcg (1.5 mL)	Oral
After 12 hours	40 mcg (2 mL)	Oral
After 14 hours	40 mcg (2 mL)	Oral
After 16 hours	40 mcg (2 mL)	Oral
After 18 hours	50 mcg (2.5 mL)	Oral
After 20 hours	50 mcg (2.5 mL)	Oral
After 22 hours	50 mcg (2.5 mL)	Oral
Once labour is established, do not increase the dose further. Maintain the current dose or decrease if tachysystole develops.		

Southern Health-Santé Sud Oral Misoprostol Titration Protocol "B" (200 mcg tablet)

Time/Progress Marker	Dose	Route
<i>Dissolve 200 mcg Misoprostol tablet in 10 mL of water = final concentration 20mcg/mL Dilute further in 30 mL of water for patient to ingest full dose</i>		
First dose after admission	20 mcg (1 mL)	Oral
After 2 hours	20 mcg (1 mL)	Oral
After 4 hours	30 mcg (1.5 mL)	Oral
After 6 hours	30 mcg (1.5 mL)	Oral
After 8 hours	40 mcg (2 mL)	Oral
After 10 hours	40 mcg (2 mL)	Oral
After 12 hours	50 mcg (2.5 mL)	Oral
After 14 hours	50 mcg (2.5 mL)	Oral
Once labour is established, do not increase the dose further. Maintain the current dose or decrease if tachysystole develops.		